


# **MME Seasonal Prediction & its Localization using CLIK**



&



**Generating  
Seasonal Outlook**

# Downscaling of seasonal prediction

Yun-Young Lee




 

**Why downscaling is required?**


# 1. Low skill of dynamical seasonal forecast...

- near surface temperature...


MAM




JJA



SON




DJF




# 1. Low skill of dynamical seasonal forecast...

- *Rainfall ?*
- *Outside tropical Pacific?*


MAM




JJA



SON




DJF



# 2. Climate Locality



## Very small Island Climate




✓ Can 2.5 by 2.5 grided model tell the difference of small island climate and ocean climat? Not really.

# Island of Sri Lanka

- *Horana in wet zone vs Angamedilla in dry zone*
- **Almost opposite temporal variation between two stations in a grid**




AMJ, cor=-0.28




# 2. Climate Locality

20 stations (Peru + Chile)



- Complicate climate of Chile




✓ Can 2.5 by 2.5 grid pixel simulate **steep terrain** effect therefore **locality of station climate** ? Not really.

# 2. Climate Locality

How different is the climate between 2 adjacent stations **in the east of Arequipa?**  
But, they are in one grid! OMG!!!


## East of Arequipa [290, -15]




# Northern arid region: RR Coast & Inland?

- 2 stations (Calama & Antofagasta) in a grid (2.5degLat by 2.5 degLon)
- Less consistency among stations in a grid


Seasonal cycle




All Chilian staions



ASO



Selected stations




**Rain shadow with strong solar insolation**



# Northern arid region: $T_{max}$ (Coast vs Inland)

- 2 stations (Calama & Antofagasta) in a grid (2.5 by 2.5)
- Less consistency among stations in a grid


Seasonal cycle




All Chilean stations




ASO: corr. coef. between stns



ASO




Selected stations




# Central region: RR Slope & Valley?

- 4 stations in a grid (2.5 degLat by 2.5 degLon)
- strong temporal consistency among stations in a grid


Seasonal cycle




All Chilean stations




ASO



Selected stations




corr. coef. between stns



# South region: RR Slope & Inland?

- 3 stations in a grid (2.5degLat by 2.5 degLon)
- Aysen: less consistent with other stations in a grid


Seasonal cycle




All Chilean stations




ASO



Selected stations



ASO: corr. coef. between stns




 

**What is downscaling?**

# Who are you?



# Look inside!






**What is downscaling?**

**POST-PROCESSING!**

**=optimization/localization/customization of  
climate information**



# What is dynamical downscaling?



- Simply, it is running a regional climate model (**RCM**).


- BC from GCM, IC  $\rightarrow$  solving dynamic equations!

- 1 month computing time for 1 month prediction ?




# Empirical-statistical downscaling

- Based on empirical relationship between **precipitation/temperature** at particular stations and **in-situ/remote large scale Atmospheric/Oceanic condition**
- Developing simple downscaling model (regression)



# Google uses AI to sharpen low-res images



# CLIK Downscaling

- ✓ *In changing climate ...*

  - : increase of climate related disasters

- ✓ *The gaps...*

  - : low dynamical prediction skill, coarse model resolution but climate locality, limited human and material resources

- ✓ *The needs...*

  - : prediction of next season climate in local community



<http://clik.apcc21.org/>

- ✓ *What we have!*

  - : Prcp/temp station data (past) & coarse GCM data (past & future)

- ✓ *What we know!*

  - : past relationship (site – large scale field)


# Approach

- Statistical downscaling forecast based on *past forecast*

$y(s, t)$  : observation

$x(s, t)$  : forecast

$$y'(t) = f(x(t), \alpha), \alpha = g(x(1 : t - 1), y(1 : t - 1))$$



- The most common way : Regression

$$\sum_j b_j y_j = \sum_i a_i x_i + \epsilon$$

If  $i$  &  $j = 1$  : Linear regression  
 $i > 1, j = 1$  : Multiple regression  
 $i$  &  $j > 1$  : CCA, SVD, etc



# Regression concept!

Scatter plot & least square fitting!

$$Y = a * x + b$$


x: predictor (ONI)

y: predictand (rainfall)



# Weakness : **overfitting**


## ■ Consider potential predictability



If model output is fitted to the unpredictable noise : Overfitting.  
What if we remove “noise” in the observation?

# Predict yield of Greek bonds with number of Facebook users

Is it appropriate?




If yes, why?

If not, why?

From *business week*

# Predict global average temperature with Carbon dioxide concentration

Is it appropriate?



If yes, why?

If not, why?

Image Source: [images.iop.org](http://images.iop.org)

# The most important things...

1. Physical understanding of,
  - What weather event/system consists of your seasonal climate (LOCAL, predictand)
  - What external (slow varying factor) controls the weather system (GLOBAL, predictor)

→ Finding **predictors** (large scale meteorological patterns (circulations) associated with local prec/temp of your station)
2. whether GCM (MME) is able to reproduce those patterns/relationship?

→ **Applicability** of downscaling



## Example of Domain Selection

:should be based on **large scale pattern** associated with local temperature/rainfall




# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)


sst vs. prec over Vientiane

JJA



sst (scm) vs. prec over Vientiane

JJA



ENSO system:

El Nino → Vientiane rain ↑

La Nina → Vientiane rain ↓


# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)


sst vs. prec over Sagaing

SON



sst (scm) vs. prec over Sagaing

SON



Indian Ocean Dipole:





Neg. IOD  $\rightarrow$  Sagaing rain  $\uparrow$

Pos. IOD  $\rightarrow$  Sagaing rain  $\downarrow$

# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center

© APEC Climate Center

IOBM:

IO warming → Dewahuwa rain ↑


IO cooling → Dewahuwa rain ↓

# Station to LSMP relationship

**OBS (Reanalysis)**

sst vs. prec over Yangon


MJJ



**Model (SCM)**

sst (scm) vs. prec over Yangon

MJJ



**Indian Ocean Basin-wide Cooling:**

SST ↓ → Yangon rain ↑

SST ↑ → Yangon rain ↓

# Station to LSMP relationship

OBS (Reanalysis)


Model (SCM)

sst vs. prec over Semtokha

DJF

sst (scm) vs. prec over Semtokha

DJF



IOBM + ENSO:





IO warming & El Nino → Semtokha rain ↑

IO cooling & La Nina → Semtokha rain ↓

# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center

© APEC Climate Center



**IOBM + ENSO:**

IO cooling & La Nina → Horana rain ↑

IO warming & La Nina → Horana rain ↓

# Maritime Continent: Indonesia

## SON [Jakarta]





**La Nina (and negative IOD) signature**



# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center



© APEC Climate Center

ENSO system:





El Nino → Osorno rain ↑

La Nina → Osorno rain ↓

# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center

© APEC Climate Center

**ENSO + IOBM:**


La Nina + Indian Ocean Basin Warming → Visviri rain ↑

El Nino + Indian Ocean Basin Cooling → Visviri rain ↓

# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center

© APEC Climate Center

ENSO + Indian Ocean Dipole :


El Nino + Pos. IOD → Santiago rain ↑

La Nina + Neg. IOD → Santiago rain ↓

# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center

© APEC Climate Center

ENSO + Tropical Atlantic Ocean Variability:


La Nina + Atlantic Nino → Visviri rain ↑

El Nino + Atlantic Nina → Visviri rain ↓

# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center

© APEC Climate Center

Non-canonical ENSO system:


CP-type El Nino → Puerto Montt rain ↑

CP-type La Nina → Puerto Montt rain ↓

# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



**Southern Tropical Atlantic Ocean Variability:**





Tropical Atlantic Cooling → Valparaiso rain ↑

Tropical Atlantic Warming → Valparaiso rain ↓

# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center

© APEC Climate Center

Rossby wave & warm easterly:



Easterly → Santiago rain ↑

Westerly → Santiago rain ↓



# Station to LSMP relationship

OBS (Reanalysis)

Model (SCM)



© APEC Climate Center




© APEC Climate Center

Large scale convective system:

Wet → Isla de Pascua rain ↑







Dry → Isla de Pascua rain ↓

# Downscaling/tailoring of dynamical MME









- 1) Physical/dynamical process
- 2) Model biases vs Observed dynamics







# Northern arid region



# Central region



# South region



# The most important things...

1. Physical understanding of,
  - What weather event/system consists of your seasonal climate (LOCAL, predictand)
  - What external (slow varying factor) controls the weather system (GLOBAL, predictor)


→ Finding *predictors* (**large scale meteorological patterns (circulations)** associated with local prec/temp of your station)
2. whether GCM (MME) is able to **reproduce those patterns/relationship?**

→ **Applicability** of downscaling

# CLIK downscaling

## ➡ A way to localize existing coarse climate information

CLIK downscaling is mainly based on station to Large Scale Meteorological Field (LSMF) relationship. ( $Y = a * X + b$ ) By utilizing the simulated LSMF (X, predictor), CLIK estimates seasonal mean precipitation/temperature (Y, predictand) at specific station.





# CLIK downscaling

## ➡ A way to localize existing coarse climate information

CLIK downscaling is mainly based on station to Large Scale Meteorological Field (LSMF) relationship. ( $Y = a*X + b$ ) By utilizing the simulated LSMF ( $X$ , predictor), CLIK estimates seasonal mean precipitation/temperature ( $Y$ , predictand) at specific station.

## Empirical relationship: LSMP (OBS) ~ local station rainfall


LSMP (MME) → Local station rainfall



# Downscaled forecast at a given site

Deterministic forecast with tercile range for target year/season



Historical time series:  
obs & downscaled from MME





Temporal Correlation Coefficient (TCC) skill

# 2018 SON Temp. forecast (Busan, KOREA)


PMME (dynamical, grid) & Downscaled (pointwise)



Busan-Gyeongsangnamdo  
Mean Temperature  
(01Sep2018 ~ 30Nov2018)



Mean	: 12.3 (Geocheon ) ~ 16.9 (Busan )
Anomaly	: -0.7 (Hapcheon ) ~ -0.3 (Ulsan )
Area Mean	: 14.5 (normal : 14.9) Anomaly : -0.4



Lead Month: 3, Year: 2018, Season: 9, Methods: GAUS

Model: APCC,CMCC,CWB,MSC,NASA,NCEP,PNU,POAMA



created by CLIK(2018-8-22)

© APEC C



# 2019 AMJ Temp. forecast (Busan, KOREA)

## PMME (dynamical, grid) & Downscaled (pointwise, SCM)



© APEC Climate Center

<b>Busan 2019 AMJ mean temp.</b>	<b>17.87 degC</b>
Normal (median)	17.27 degC
	<b>+ 0.6 degC</b>



# **Availability of Downscaled information?**



# PICASO

Pacific Island Countries Advanced Seasonal Outlook

# Downscaling/tailoring of dynamical MME



Dynamical Prediction

Current Climate



They further tailor “predicted climate”  
onto  
“predicted rainfall” at the given site.

Predicted Climate

Predicted Rainfall

- 1) Physical/dynamical process
- 2) Model biases vs Observed dynamics

# ROK-PI CliPS: Overview

Republic Of Korea-Pacific Islands Climate Prediction Services

- Objective:

*“To strengthen the adaptive capacity of vulnerable communities to climate risks at the seasonal timescale.”*

- Working Pillars:

**One** Dynamical Seasonal Forecasting System



**Two** Downscaled Prediction System



**Three** Development of the Application Guideline



**Four** Training of the Climate Information Application



# Intuitive climate outlook





Figure 2. Temporal correlation coefficients (TCCs) between local precipitation of Butaritari(91601) and precipitation at each grid during the April – June(AMJ) for (left) observation and (right) DMME. The black dots indicate grid points for which TCC is significant at the 95% confidence level.

The large-scale oceanic and atmospheric signals associated with the local precipitation at Butaritari(91601) during April-June(AMJ) season are displayed in Figure 1 and 2. The dynamical seasonal prediction system (APCC-MME based PICASO) represents that the AMJ precipitation of Butaritari(91601) is wellrelated to warm equatorial Pacific state (e.g., El Niño), and it can be best recognized by the predicted (MME) remoteprecipitation over the Western Pacific. Therefore, remoteprecipitation is selected as the internal predictor in PICASO.

**TRAINING SCORES**

Name	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ	TOTAL
HSS	0.14	0.2	0.14	0.57	0.5	0.03	0	0.32	0.14	0.28	0.09	0.11	0.21
LEPS	33%	35%	27%	31%	35%	22%	35%	20%	22%	38%	38%	23%	0%

**TRAINING HISTORY (PREDICTION/OBSERVATION)**

Name	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
2005	0/0	0/-	+0	0/+	-/+	+/+	0/+	+0	0/+	+/-	0/+	+/+
2004	0/+	0/+	0/0	0/0	0/0	+0	0/+	0/0	0/0	0/+	0/+	+/-
2003	0/0	+/-	0/0	0/+0	-/+	0/+0	+0	+0	+0	+0	+0	0/+
2002	0/0	0/+0	0/0	0/0	+/-	-/+	0/0	0/+0	0/+0	+0	+0	0/+
2001	+/-	0/+0	0/0	0/+	0/+0	-/+	0/+0	0/0	0/+	0/+0	+0	0/+
2000	+/-	0/0	+/-	+/-	0/+0	+/-	0/0	-/+	-/+	-/+	+0	0/+
1999	0/+0	0/+	+/-	0/0	0/0	0/0	+0	+0	+0	+0	0/+0	+0
1998	0/0	0/+0	0/0	0/0	-/+	0/0	0/+0	0/+0	0/0	0/0	0/0	0/+
1997	0/+	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/+
1996	0/+	0/+0	0/+	-/+	0/+0	0/+	0/+0	0/+	0/0	0/+0	0/+	+0
1995	0/+	0/0	-/+	0/0	0/+	0/0	0/+	0/0	0/0	+0	0/+	+0
1994	0/0	+/-	+/-	0/0	0/0	0/+	+0	0/0	0/+	0/0	0/0	+0

**VALIDATION SCORES**

Name	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ	TOTAL
HSS	0.14	0.2	0.14	0.57	0.5	0.03	0	0.32	0.14	0.28	0.09	0.11	0.21
LEPS	33%	35%	27%	31%	35%	22%	35%	20%	22%	38%	38%	23%	0%


**VALIDATION HISTORY (PREDICTION/OBSERVATION)**

Name	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
2016	0/0	0/-	+0	0/+	-/+	+0	+0	+0	+0	+0	+0	+0
2015	0/+	0/+	0/0	0/0	0/0	0/+	0/+	0/+	0/+	0/+	0/+	0/+
2014	0/0	+/-	0/0	0/+0	+/-	0/+0	+0	+0	+0	+0	+0	+0
2013	0/0	0/+0	0/0	0/0	+/-	-/+	0/0	0/0	0/0	0/0	+0	+0
2012	+/-	0/+0	0/0	0/+	0/+0	-/+	0/+0	0/0	0/+	0/+0	0/+0	0/+0
2011	+/-	0/+	+/-	+/-	0/0	+0	0/0	0/0	0/0	0/0	+0	+0
2010	0/+0	0/+	+0	+0	0/0	0/+	0/+0	0/+0	0/+0	0/+0	0/+0	0/+0
2009	0/0	0/0	0/0	0/0	-/+	0/0	0/+0	0/+0	0/0	0/0	0/0	0/+
2008	0/0	0/0	0/0	0/0	-/+	0/0	0/+	0/+	0/+	0/+	0/+	0/+
2007	0/+	0/+0	0/+	-/+	0/+0	0/+	0/+0	0/+	0/0	0/+0	0/+0	0/+0
2006	0/+	0/0	-/+	-/+	0/0	0/+	0/0	0/+	0/0	0/+	0/+	0/+

# PICASO > Details > Probability Distribution

- **Interactive Probability Scale**

*Probability below/above specific criteria*



**Chance of rainfall more than  
299.3mm/mon during SON, 2017  
= 50.9%**

**Chance of rainfall less than  
299.3mm/mon during SON, 2017  
= 49.1%**

- **Applicable to other sectors**

Water  
resources  
management

Disaster Risk  
Reduction


Agricultural  
planning

# Tuvalu

## Current Water Resources

- **Primary - Rain water (rain harvest and banked at individual household)**
- Supplementary - RO desalination plant source from either the groundwater or direct from the sea.
- Secondary - ground water (only for secondary usage - lavatory cleaning, bathing) mainly during the dry spells.

Average annual rainfall: 3000mm ~ 4000mm (in the southern islands)



**WARNING!**

# Samoa

## Watershed Management Plans

### Zoning regime

*Zone 3: Exclusion Zone (above 600m altitude)*

*Zone 2: Restricted Zone (300m-600m above mean sea level)*


*Zone 1: Development Zone (0-300m above mean sea level)*

- ✓ Zone 1 is the lower watershed area where **population/development** is concentrated and where the **needs for watershed ecological services** are greatest.
- ✓ Development planning is essential in implementing **flood hazard plans** and also in ensuring that activities have controlled impacts on the receiving coastal waters and soil resources.


### Abundant water resources

- The Average rainfall is over 3,000mm/year (2,500 mm in the north-west parts of the main islands to 6,000 mm in the highlands of Savaii)
- 75 percent of the precipitation occurring during November-January.

# NO WORRIES!



# Kiribati




**WARNING!**

- A vast area of territory covering most equatorial Pacific
- Regionally/seasonally varying - ENSO dependent rainfall
- La Nina coming → Drought expected  
→ Advise **the Drought Committee** on the status of the Water reserve

# Tonga

## Recharge to groundwater



Annual Rainfall: 1,770 mm

Potential Evaporation: 1,550 mm

Tonga Water Board (TWB)  
: **water supply services** for domestic, stock, horticultural, industrial, commercial, recreational, environmental and other beneficial uses

## TONGA'S FIRST COMMERCIAL CONSIGNMENT OF SQUASH PUMPKIN IS ON ITS WAY TO CHINA



(Radio & TV Tonga, Nuku'alofa, 28/11/2016)

The Chinese Government has approved the export of Tonga's squash pumpkin into its market.

This is seen as an initiative that could encourage more growers to go back to the plantations they abandoned during the past years due to falling prices in overseas markets.


The CEO of Agriculture – Dr. Viliami Manu told Radio Tonga news, this is a result of an evaluation visit of a team from China's Quarantine Department to oversee the quality of squash pumpkin in Tonga.

<http://www.tonga-broadcasting.net/?p=3959>


### Watering

For good growth, squash and pumpkins require at least one inch of water per week. (One inch of water per thousand square feet is 620 gallons). If water is needed, irrigate thoroughly early in the morning until the soil is moistened eight to twelve inches deep. If rainfall is deficient, it may be necessary to water once a week, perhaps two times per week in sandy soils.

[https://ag.umass.edu/sites/ag.umass.edu/files/fact-sheets/pdf/pumpkins\\_and\\_squash.pdf](https://ag.umass.edu/sites/ag.umass.edu/files/fact-sheets/pdf/pumpkins_and_squash.pdf)




# Fill the gap



**Accuracy**  
**Resolution**  
**Contents**







# Climate Prediction Service Enhancement












Thank you.

# Tx



# Tn



# RR

